



Full Length Research Article

A COMPARATIVE STUDIES OF TWO SILKWORM SPECIES IN RAIGARH DISTRICT FOR HIGH QUALITY YIELD

*Sheela Patel, R. K. Singhand Shweta Sao

Department of Life Sciences, Dr. C.V. Raman University, Bilaspur, Chhattisgarh, 495113, India

ARTICLE INFOABSTRACT

Article History:

Received 27th September, 2016
Received in revised form
22nd October, 2016
Accepted 19th November, 2016
Published online 30th December, 2016

Key Words:

Sericulture, Silkworms,
Antheraea mylitta, *Bombyx mori*,
Comparative studies.

The sericulture industry is very unique and popular for its high valued textile fiber, agricultural outputs including, cocoons and cottage based labor intensive and for the production of high quality textiles. An attempt has been made to formulate a strategic model to recognize the potential strength and challenges of sericulture in Raigarh district at Chhattisgarh state. We report results of the comparative studies relating to the silk production and productivity at Raigarh district from the silkworm species *Antheraea mylitta* and *Bombyx mori*. The study indicates the commercial perspective of silkworm production at Raigarh district. The shell weight of *Antheraea mylitta* and *Bombyx mori* (female species) obtained as 0.95 ± 0.06 and 0.26 ± 0.01 respectively, which confirms the higher production and yield as compared to *Bombyx mori*.

Copyright©2016, Sheela Patel et al., This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

India continues to be the second largest producer of silk in the world, providing gainful occupation to about six million persons in the rural and semi-urban areas across the country (Rahaman, 2013). India has unique distinction of being the only country on the world producing all the four commercially known varieties of silk, viz, mulberry, tasar, eri and munga (Anitha et al., 2011). Sericulture industry has been identified as employment oriented industry (Devaraja, 2011). All the sections of sericulture industry, viz. mulberry cultivation, silkworm seed production, silkworm rearing, reeling and weaving of silk and collection of byproducts and its processing provide a large scale employment, thereby a source of livelihood for the rural and tribal people (Gregory, 1994 and Srivastava et al., 2003). Sericulture industry is rated as the second largest employer in India (Anitha, 2011). Sericulture have recently aroused much interest due to their wide industrial applications. Silk is the most elegant textile in the world with unparalleled grandeur, natural sheen, and inherent affinity for dyes, high absorbance, light weight, soft touch and high durability and known as the "Queen of Textiles" the world over (Li et al., 2012; Bunning et al., 1994). Silk is composed of two major proteins, a fibrous protein (fibroin) and globular, gumming protein such as sericin (Takasu et al., 2002; Mondal, 2007).

Fibroin has been used in textile manufacturing and for several biomaterial and biomedical applications, whereas sericin is considered a waste material in the textile industry (Reddy and Prasad 2011; Vepari and Kaplan 2007). Sericin has recently been found to activate the proliferation of several cell-lines and has also shown various biological activities (Tsubouchiet al., 2005; Zhang et al., 2006). Chhattisgarh state is a very popular for its high quality Kosa silk production. Silk is a way of life in Chhattisgarh has become an inseparable part of Indian culture and tradition should be considered for rural management and development (Dewangan et al., 2011). Presently in Chhattisgarh three types of silk viz., Mulberry, Tasar and Eri silk are producing (Singh, 1993). Mulberry and Tasar farming is done in 6 major district such as Raigarh, Bilaspur, Korba, Champa, Bastar and Ambikapur.

Chhattisgarh is the second largest producer of Tasar cocoons. Mainly two types of silk produced in the Raigarh district; they are tasar silk & mulberry silk. Kosa of Chhattisgarh was famous in the world for its elegance and quality and it is still, a force considers. Raigarh district is major tasar growing area where tribal are engaged in sericulture activities (Dewangan et al., 2011). The Kosa industry is one of the industries that is currently blooming in Raigarh along with other industries producing iron ore, steel and power in the country. The production of Kosa is the main livelihood for some of the villagers in this district and many of them have now started running units for producing Kosa sarees

*Corresponding author: Sheela Patel, Department of Life Sciences, Dr. C.V. Raman University, Bilaspur, Chhattisgarh, 495113, India.

and dress materials for export (Dewangan et al., 2011). The present investigation was carried out in Raigarh district of Chhattisgarh state. Here, we report the results of the studies relating to comparative study of silk production at Raigarh district. Present study focused on two silkworm species *Antheraea mylitta* and *Bombyx mori*. Specifically, the main objectives of the study are: (1) to introduce in brief, the historical perspective of sericulture in Raigarh district at Chhattisgarh state. (2) Comparative Studies of two silkworm species such as *Bombyx mori* and *Antheraea mylitta* and their life cycles studies. (3) To examine the growth and production of sericulture in Raigarh district at Chhattisgarh state (4) Study of the improvements of silk quality and production of silk up to industrial scale.

MATERIALS AND METHODS

Field of study

The present research study has been carried out in Raigarh district at Chhattisgarh state (Figure 1). Raigarh is the second largest producer of tasar cocoons in Chhattisgarh state, where both types of silk namely mulberry and tasar are being produced. Based on kosa silk, many villagers are running units for producing kosa sarees and dress materials of export quality. The study area has about 364 acres under mulberry cultivation though effective area is only 80-120 acres. Mulberry gardens are 18 in number and Mulberry Grainages are 1 in Urdana.

Mulberry Reeling Unit in 1 number at Dharamjaigarh. Raigarh district is major tasar growing area where tribal are engaged in Sericulture activities. Tasar culture is a traditional and exclusive craft of the tribal of study area and is being practiced from 30 years. Maximum number of total production of tasar cocoons in the State comes from the district. Total area covered under tasar centers is about 3153.25 acres though effective area is about 2350 acre. Tasar centers are 22 in number (Dewangan et al., 2011).

Climatic factors

The weather in Raigarh varies during different months. In the summer the temperature can vary from 29.5 °C to 49 °C and lasts during the month from March to May. The temperature during this pre-monsoon period increases rapidly during the night and day.

It's usually extremely dry during this time. During the monsoon which last from June to September, the maximum temperature is 38 °C in the month of June and in the month of September the temperature goes down to 24.50 °C. At times, the monsoon also extends up to mid-October. Even though the temperature drops to 17.10 °C during the nights of October, the official winter season starts from December and goes on to the month of February to January with the minimum temperature going down to 13.20 °C (Chakraborty et al., 2015).




Figure 1. Sericulture in Raigarh district at Chhattisgarh state

Sample collections

The present investigation was carried out in Raigarh district at Chhattisgarh state, was intentionally selected for the study, as it is the second largest producer of Tasar cocoons in Chhattisgarh, where both types of sericulture - mulberry and tasar are being accomplished. The present research study has been carried out in nearby sericulture research centre in Raigarh district such as Baramkela, Boirdadar and Deacon.

Rearing of silkworm

Rearing of Tasar and mulberry silkworms was carried out on Arjun (*Terminalia arjuna*) and mulberry plant respectively. For this, freshly hatched healthy silkworms were brushed on to each of the plant species during the 1st crop (July - August) season following the rearing protocol developed by Central Silk Board, Bangalore (Dandin and Giridhar, 2010).

Mulberry silkworm

Cultivation of mulberry plant is mainly for its leaves the sole food for the silkworm, *Bombyx mori* L. for commercial production of raw silk. The silkworms are actually larvae of the silk moths shown in Figure 2A and Figure 2B. They are reared in specially made trays in rooms with controlled temperature and humidity and regularly fed mulberry leaves. At a certain stage they convert themselves into cocoons. These cocoons are made from a single filament of material secreted by the pupa and wrapped around itself for protection. These filaments upon hardening constitute silk. On an average, 1 acre of plantation would yield 240 kg of cocoons in a year, starting from 100 DFLs. Depending upon whether it is dryland or irrigated mulberry, farmers can harvest the cocoons 4 to 8 times in a year.

Tasar silkworm

The tasar silkworms belong to the genus *Antheraea* and they are all wild silkworms. The Indian tasar worms feeds on leaves of *Terminalia* and several other minor host plants. The *Antheraea mylitta* silkworms larvae and moth as shown in Figure 2C and Figure 2D. The worms are either uni or bivoltine and their cocoons like the mulberry silkworm cocoons can be reeled into raw silk. In Tasar culture, the silkworms are reared outdoors on the trees, for better management of rearing it is desirable that the plants are given proper height and shape. Since in Tasar culture, it is the leaf and not the wood or fruits, which are required, plants should be induced to produce more of quality leaves. The plants are cut at 6' height for Arjun and 7' height for Asan while in light pruning only the branches of 1' diameter or less are cut. Arjun and Asan plantation of more than four years ago with spacing of 4x4' are maintained by pruning at the height of 3-4'. pruning should be done during February-March and march-April for 1st and 2nd crop, respectively.

Life cycle study

Life cycle of *Antheraea mylitta* and *Bombyx mori* have been studied for comparative studies. Life cycle of the silkworm consists of four stages i.e. adult, egg, larva, and pupa. The duration of life cycle is six to eight weeks depending upon racial characteristics and climatic conditions. Multi-voltine races found in tropical areas have the shortest life cycle with

the egg, larval, pupal and adult stages lasting for 9-12 days, 20-24 days, 10-12 days and 3-6 days, respectively.




Figure 2. (A) *Bombyx mori* larva, (B) moth on mulberry plant, (C) *Antheraea mylitta* larva and (D) moth on arjun plant respectively.




Figure 3. Life cycle of silkworm *Bombyx mori*




Figure 4. Life cycle of silkworm *Antheraea mylitta*

Seven to eight generations are produced in multi-voltine races. In uni-voltine races, the egg period of activated egg may last for 11-14 days; the larval period, 24-28 days; the pupal period, 12-15 days and the adult stage, 6-10 days. The life cycle of *Bombyx mori* has been shown in Figure 3. In nature, uni-voltine races produce only one generation during the spring and the second generation of eggs goes through a period of rest or hibernation till the next spring. In case of bivoltine races, however, the second generation eggs do not hibernate and hatch within 11-12 days and produce second generation normally during summer and it is the third generation eggs which undergo hibernation and hatches in the next spring, and thus producing two generations in one year. Several species of *Antheraea* are exploited for production of wild silk known as tasar silk. The tasar moths are fairly large insects. Females are larger and yellowish brown in colour, while males are smaller and brick red in colour. Both have prominent and colourful eye spots on their wings. The antennae of males are bushy, and abdomen is narrower in comparison to female. The important stages of the tasar silkworm life cycle have shown in Figure 4.

status of host plant but farmers for the economic advantage use alternative food plants based on availability and accessibility. The rate of leaf production, quantity, gestation period of host plants in relation to season compared to primary food plant and their commercial feasibility is an important factor for silkworm rearing and grainage performance. The present study shows that tasar rearing and grainage behaviour is better when the larvae were fed *Terminalia arjuna* food plant though commercial traits viz., cocoon weight, shell weight, silk ratio and egg fertility are much better than mulberry host plant for *Bombyx mori*. This is indicative of availability of suitable nutrients in the host plant *Terminalia arjuna*. The highest cocoon weight 12.35 gm was recorded for species *Antheraea mylitta* (female) as compared to *Bombyx mori* (female) cocoon weight 1.47 gm rainy season which are shown in Table 1 and Table 3, respectively. Wherever, the highest cocoon weight 9.30 gm was recorded for species *Antheraea mylitta* (male) as compared to *Bombyx mori* (male) cocoon weight 1.23 gm rainy season which are shown in Table 2 and Table 4, respectively.




Figure 5. (A) Comparative statistical graph of cocoon parameter of *Antheraea mylitta* and *Bombyx mori*(male), (B) Comparative statistical graph of cocoon parameter of *Antheraea mylitta* and *Bombyx mori* (female).

Statistical analysis

Research methodology applied in this study was a combination of descriptive-analytical and quantitative methods and statistical methods. Primary and secondary data was analyzed using various statistical tools viz., mean, mode and median. In addition to usual statistical measures coefficient correlation techniques were employed at appropriate context in the study to evaluate and analyze the collected data.

RESULTS AND DISCUSSION

The results in this study has been supported by the statistical data analysis presented in the form of tables given under different sections. Cocoon yield and seed quality of *Bombyx mori* and *A. mylitta* is dependent on variety and nutritional

Productivity of cocoon in outdoor rearing is poor due to attack of number of pests and predators besides natural vagaries such as, wide fluctuating temperature, heavy rain and stormy wind etc. Attempts were made in the past to increase the cocoon production by adopting various methods of indoor rearing was conducted in wooden tray supported with wooden rearing frames specially designed for young age and late age silkworms. Production of quality tasar seed is one of the most challenging task of tasar silk industry at base level. A systematic and methodological approach of silkworm seedproduction is required to sort out the problems during the preparation of quality seed. Tropical tasar *Antheraea mylitta* is production is required to sort out the problems during the preparation of quality seed. Tropical tasar *Antheraea mylitta* is wild sericigenous insect of commercial importance of tropical India.

Table 1. Data showing individual length (cm), shell and pupa weight (gm), shell weight (gm) and pupa weight (gm) of *Antheraea mylitta*(female)

S.N.	Length (cm) [A]	Shell +pupa weight (gm) [B]	Shell weight (gm) [C]	Pupa weight (gm) [D]
1.	4.50	8.34	1.05	7.29
2.	4.80	9.09	1.23	7.86
3.	4.70	9.30	1.28	8.02
4.	4.40	7.91	0.91	7.00
5.	4.00	7.55	0.93	6.62
$\sum x/N$	22.4/5	42.19/5	5.4/5	36.79/5
Mean	4.48	8.44	1.08	7.35
SD	± 0.31	± 0.75	± 0.17	± 0.59
SE	± 0.14	± 0.34	± 0.07	± 0.26

Table 2. Data showing individual length (cm), shell + pupa weight (gm), shell weight (gm) and pupa weight (gm) of *Antheraea mylitta*(male)

S.N.	Length (cm) [A]	Shell and Pupa weight (gm) [B]	Shell weight (gm) [C]	Pupa weight (gm) [D]
1.	5.20	12.30	0.97	11.33
2.	4.50	12.35	1.11	11.24
3.	5.00	11.37	1.05	10.32
4.	5.10	9.46	0.81	8.65
5.	4.80	8.54	0.83	7.71
$\sum x/N$	24.6/5	54.02/5	4.77/5	49.25/5
Mean	4.92	10.80	0.95	9.85
SD	± 0.28	± 1.72	± 0.13	± 1.61
SE	± 0.12	± 0.77	± 0.06	± 0.72

Table 3. Data showing individual length (cm), shell + pupa weight (gm), shell weight (gm) and pupa weight (gm) of *Bombyx mori*(female)

S.N.	Length (cm) [A]	Shell and Pupaweight (gm) [B]	Shell weight (gm) [C]	Pupa weight (gm) [D]
1.	3.70	1.46	0.31	1.15
2.	3.50	1.47	0.25	1.22
3.	3.00	1.33	0.25	1.09
4.	3.30	1.06	0.22	0.84
5.	3.20	1.45	0.25	1.20
$\sum x/N$	16.7/5	6.77/5	1.28/5	5.49/5
Mean	3.34	1.35	0.26	1.09
SD	± 0.27	± 0.17	± 0.03	± 0.15
SE	± 0.12	± 0.08	± 0.01	± 0.07

Table 4. Data showing individual length (cm), shell and pupa weight (gm), shell weight (gm) and pupa weight (gm) of *Bombyx mori* (male)

S.N.	Length (cm) [A]	Shell and Pupaweight (gm) [B]	Shell weight (gm) [C]	Pupa weight (gm) [D]
1.	2.60	0.89	0.25	0.64
2.	2.90	0.69	0.22	0.47
3.	3.00	1.23	0.21	1.02
4.	2.90	1.05	0.22	0.83
5.	3.00	0.63	0.28	0.35
$\sum x/N$	14.4/5	4.49/5	1.18/5	3.31/5
Mean	2.88	0.89	0.24	0.66
SD	± 0.16	± 0.25	± 0.03	± 0.27
SE	± 0.07	± 0.11	± 0.01	± 0.12

Table 5. Achievements of Basic Seed Multiplication and Training Centre, Boirdadar, Raigarh in cocoon production from year 2006 to 2016.

S.N.	Years	Protege Healthy Group	Cocoon Production
1.	2006-2007	18500	1019264
2.	2007-2008	21500	1154525
3.	2008-2009	27000	1390885
4.	2009-2010	23375	1571246
5.	2010-2011	25100	1749295
6.	2011-2012	27800	2005475
7.	2012-2013	25885	1860470
8.	2013-2014	31000	1858970
9.	2014-2015	32275	1793050
10.	2015-2016	34290	1997062

The cocoon traits like cocoon length and cocoon weight, single shell weight, pupal weight in rainy seasons is related to the host plants fed to larvae. The comparative statistical data of cocoon parameter from *A. mylitta* and *Bombyx mori* male species have shown in Figure 5A. The shell weight of female *Antheraea mylitta* is $0.95 \pm 0.06^*$ as compare to female *Bombyx mori* shell weight 0.26 ± 0.01 . Similarly, the male *Antheraea mylitta* shell weight is $1.08 \pm 0.07^*$ as compared to male *Bombyx mori* shell weight 0.24 ± 0.01 , which confirms the higher productivity of *Antheraea mylitta* as compared to *Bombyx mori*.

The comparative statistical data of cocoon parameter from *A. mylitta* and *Bombyx mori* female species have shown in Figure 5B. Present study confirms the high quality production and yield obtained from *Antheraea mylitta* silkworm in Raigarh district. The cocoon traits like cocoon length and cocoon weight, single shell weight, pupal weight in Rainy seasons is related to the host plants fed to larvae. Multiplication and Training Centre, Boirdadar, Raigarh have shown in have shown in Figure 6. Target and achievements of Achievements of Basic Seed Multiplication and Training Centre (BSMTC), Boirdadar, Raigarh in cocoon production from year 2006 to 2016 has been mentioned in Table 5. The data shows that the large amount of cocoon production in the year 2016 up to 1997062 in farming area 117.81 hectare.

Conclusions

Sericulture is the science that deals with the production of silk by rearing of silkworm. Sericulture plays a vital role in the flow of income from the urban rich sections of the society to the rural poor, as demand for silk is largely from the higher income group. In Chhattisgarh tropical Tasar and mulberry are reared on commercial scale. Tasar is really named as Kosa. Raigarh district is major tasar growing area where tribal are engaged in sericulture activities. Present study confirms that, bivoltine silkworms are superior over crossbreed and multivoltine silkworms in biochemical contents in different body tissues analyzed. Difference between silk worms is due to genetic endowment of the races. Quantity and quality of bimolecular in silkworms attributes the robustness and healthiness that reliably considered being better in rearing performance and cocoon yield. Screening of silkworm genetic resources using biochemical analysis as a tool may be more dependable for the selection in silkworm breeding programs as well as for commercial exploitation of silkworm races.

Acknowledgement

The authors thanks to Mr. Uday Narayan Singh, Senior Scientist, Mr. Naresh Kumar Sodiwala and Bhagirathi Patel, Basic Seed Multiplication & Training Centre, Boirdadar, Raigarh for proving the research facilities to carry out the present experimental work. Authors also thanks to Mr. T. L. Patel, Field Officer, State Govt. Mr. Sulkamber Pradhan, Senior Field Officer, Tasar Reeling Factory, Urdana, Raigarh for valuable discussions and timely support from the beginning of research work.

REFERENCES

- Anitha, R. 2011. Indian silk industry in the global scenario. International Journal of Multidisciplinary Management Studies, 1: 100-110.
- Bunning, T. J., Jiang, H., Adams, W.W., Crane, R. L., Farmer, B., Kaplan, D. 1994. Applications of silk. In ACS Symposium Series. American Chemical Society, 544:353-353.
- Chakraborty, S., Muthulakshmi, M., Vardhini, D., Jayaprakash, P., Nagaraju, J., Arunkumar, K. P. 2015. Genetic analysis of Indian tasar silkworm (*Antheraea mylitta*) populations. Sci. Rep. 5:15728.
- Dandin, S. B., Giridhar K. 2010. Handbook of Sericulture Technologies. Central Silk Board, Ed.
- Devaraja, T. S. 2011. Indian textile and garment industry-An overview. New Delhi: Indian Council of Social Science Research.
- Dewangan, S.K., Sahu, K.R., Achari, K. V., Soni, S. 2011. Socio-economic empowerment of tribal women through sericulture a study of Lailunga block of Raigarh District, Chhattisgarh, India. Int. J.B. Manag. 6: 297.
- Gregory, S. 1994. Rural Labour and Sericulture: Typology, Strategies and Prospects. Indian J. Ind. Relat. 1: 365-376.
- Li, G., Liu, H., Li, T., Wang, J. 2012. Surface modification and functionalization of silk fibroin fibers/fabric toward high performance applications. Mater. Sci. Eng. C. 32: 627-636.
- Mondal, M. 2007. The silk proteins, sericin and fibroin in silkworm, *Bombyx mori* Linn., A review. Caspian Journal of Environmental Sciences, 5:63-76.
- Rahaman, A. H., Khan, N., Khan, S. A. 2013. Spatio-temporal changes of growth and production of sericulture in Asia: An analytical and comparative study. J. Geogr. Reg. Plann. 6: 63.
- Reddy, R. M., Prasad, G. V. 2011. Silk-the prospective and compatible bio-material for advanced functional applications. Trends Appl. Sci. Res. 6: 89.
- Singh, C. 1993. An economic analysis of sericulture production in Raigarh district of Madhya Pradesh. Agricultural Economics Research Review, 6: 52-53.
- Srivastava, S., Kapoor, R., Thathola, A., Srivastava, R. P. 2003. Mulberry (*Moms alba*) leaves as human food: a new dimension of sericulture. Int. J. Food Sci. Nutr. 54: 411-416.
- Takasu, Y., Yamada, H., Tsubouchi, K. 2002. Isolation of three main sericin components from the cocoon of the silkworm, *Bombyx mori*. Biosci. Biotechnol. Biochem. 66:2715-2718.
- Tsubouchi, K., Igarashi, Y., Takasu, Y., Yamada, H. 2005. Sericin enhances attachment of cultured human skin fibroblasts. Bioscience, Biotechnology, And Biochemistry, 69:403-405.
- Vepari, C., Kaplan, D.L. 2007. Silk as a biomaterial, Prog. Polym. Sci. 32: 991-1007.
- Zhang, Y. Q., Ma, Y., Xia, Y.Y., Shen, W.D., Mao, J.P., Xue, R.Y. 2006. Silk sericin-insulin bioconjugates: Synthesis, characterization and biological activity. J. Control. Release, 115: 307-315.
