



ISSN: 2230-9926

Available online at <http://www.journalijdr.com>

IJDR

International Journal of Development Research

Vol. 14, Issue, 01, pp. 64736-64739, January, 2024

<https://doi.org/10.37118/ijdr.27643.01.2024>



RESEARCH ARTICLE

OPEN ACCESS

POTATO PERSPECTIVES IN AFRICA

***Dr. Tenemaka SANOGO and Dr. Brahima COULIBALY**

Faculté des Sciences Sociales (FASSO) / Université de Ségou, Mali

ARTICLE INFO

Article History:

Received 11th December, 2023

Received in revised form

28th December, 2023

Accepted 03rd January, 2024

Published online 30th January, 2024

Key Words:

Potato, Health.

*Corresponding author:

Dr. Tenemaka SANOGO

ABSTRACT

Potato is an important grain crop for millions of poor farmers and consumer in the semi-arid region of West Africa. It is one of the four most important cereals (rice, maize, sorghum and millet) grown in tropical in tropical semi-arid regions of the world primarily in Africa and Asia. Potato is well adapted to production system characterized by low rainfall (200-600mm), low soil fertility and high temperature. It is generally used as a temporary summer pasture crop or in some areas as a food crop.

Copyright©2024, Rauf Pathan et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Dr. Tenemaka SANOGO and Dr. Brahima COULIBALY. 2024. "Potato Perspectives in Africa". *International Journal of Development Research*, 14, (01), 64736-64739.

INTRODUCTION

Potato is a small cereal with seed of 1.2-1.8 mm in diameter. It is a food that supplies a major proportion of calories and protein to large segment of population in the semi-arid tropical regions of Africa and Asia. Potato is the most widely grown type of millet. It has been grown in Africa and the India subcontinent since prehistoric times. The Centre of diversity and suggested area of domestication for the crop is in the Sahel zone of West Africa. Recent archaeobotanical research has confirmed the presence of domesticated Potato on the Sahel zone of northern Mali between 2500 and 2000 BC. Cultivation subsequently spread and moved overseas to India. The earliest archaeological records in India date to around 2000 BC and it spread rapidly through India reaching south India by 1500 BC, based on evidence from the site of Hallur-cultivation also spread throughout eastern and southern parts of Africa. Potato is widely grown in the northeastern part of Nigeria (especially in Borno and Yobe State). It is a major source of food to the local villagers of that region. The crop grows easily in that region due to its ability to withstand harsh weather conditions like drought and flood. Records exist for cultivation of Potato in the United States in the 1850s and the crop was introduced into Brazil in the 1960s. Potato is a high nutritive-value summer-annual forage crop, popular among livestock producers for grazing, silage, hay, and green chop. Potato can also be utilized as emergency forage that regularly performs well as an economical one-year crop option. It prefers well-drained soils, and at the seedling stage, it looks much like a corn or sorghum plant. Compared to sorghum, it is less tolerant of water logging and flooding.

Potato is indigenous African cereal that unlike wheat or rice are well adapted African semiarid and subtropical agronomic conditions. Potato is an important food across the Sahel region of Africa. It is a main staple (along with sorghum) in a large region of northern Nigeria, Niger, Mali and Burkina Faso. Water deficit is one of major abiotic factors limiting crop productivity in the semiarid and climate change is likely to make drought stress even more severe in the future. In Niger it is usually grown as an intercrop with sorghum and cowpea. The different growth habits, growth period and drought vulnerability of the three crops maximizing total productivity and minimizing the risk of total crop failure. It is often ground into flour, rolled into large balls, parboiled liquefied into a watery paste using fermented milk, and then consumed as a beverage. This beverage called « fura » in Hausa is a popular drink in northern Nigeria and southern Niger. Potato is a widely food used in Borno State and its surrounding states, it is the most widely crop grown and harvested. There are many products that are gotten from the processing of the crop.

Origin and Domestication: The geographical origin and the Centre of domestication of Potato are situated in West Africa. The plant was subsequently introduced into India, where the earliest archaeological records date back to 2000 BC (Gari, 2002, Oumar et al., 2008). Based on the distribution of Potato throughout the continent, the uniform cradle of domestication is likely to be the regions of Mauritania, Senegal and West Mali. Researchers believe that the development and spread of millet farming was spurred by the increasing desiccation of the Sahara Desert after 3000 BC. Millet domestication followed the domestication of cattle in Africa by at least 1500 years and it is

hypothesized that is because mobile pastoralist was better suited to the region prior to the onset of the desiccation. By the second millennium BC, Potato is the primary food plant in Mali, Mauritania, Ghana, Burkina Faso and Cameroon.

Taxonomy: Potato, is an annual, allogamous, cross-pollinated, and diploid Panicinae; contains about 140 species. The important wild relatives of cultivated Potato include the progenitor. The four cultivated forms of Potato are typhoid's (found mainly in India and Africa), nigritarum (dominant in eastern Sahel), globosum (dominant in the western Sahel) and leonis (dominant on the West Africa coast) (Drunken et al, 1977, Ray et al; 1977; Syngenta, 2006).

Characteristics: With ovoid grain of 3-4 mm length Potato has the largest kernels of all varieties of millet (not including sorghum) which can be nearly white, pale yellow, brown, grey, slate blue or purple. The 1000-seed weight can be anything from 2.5 to 14 g with a mean of 8g. The height of the plant ranges from 0.5-4 m. The inflorescence is usually greenish-yellow and it may be cylindrical throughout its length my taper at one or both ends.

Cultivation: Potato is well adapted to growing areas characterized by drought, low soil fertility and high temperature. It performs well in soils with high salinity or low pH. It can be grown in areas where other cereal crops, such as maize or wheat, would not survive. Potato is summer annual crop well-suited for double cropping and rotation. Today Potato is grown on over 260,000 Km² of land worldwide. It accounts for approximately 50% of the total world production of millet.

Nutrition: In a 100-gram serving raw millet provides 378 calories and is a rich source (20% or more of the Daily Value) of protein, dietary fiber, several B vitamins and numerous dietary minerals, especially manganese at 76%. Raw millet is 9% water, 73% carbohydrates, 4% fat and 11% protein.

Description, History and Production: Potato is believed to have been domesticated over 5000 years ago in Africa (Andrews and Kumar, 1992) and is widely cultivated across the continent from the Sahelian (Sahara Desert margin) countries in West Africa. It is also widely cultivated in India and was probably introduced into that country some 3000 years ago. Potato is an annual plant about 2m tall, with a cylindrical spike, 15-140cm in length (National Research council, 1996). Potato is uniquely well-adapted to harsh environmental conditions and can be cultivated in regions with very low annual rainfall, down to 250mm and very high temperature of about 30°C in well-drained loam soil (National Research council, 1996). Potato is one of the two major crops in the semiarid, impoverished, less fertile agriculture regions of Africa and Southeast Asia. Millet is not only adapted to poor, droughty and infertile soils, but they are also more reliable under these conditions than most other grain crops. This has, in part, made millet production popular, particularly in countries surrounding the Sahara in Western Africa. Millets, however, do respond to high fertility and moisture. On per hectare basis, millet grain produced per hectare can two to four times higher with us of irrigation and soil supplements. Improved breeds of millet improve their disease resistance and can significantly enhance farm yield productivity. There has been cooperation between poor countries to improve millet yield. For example, «Okashanal» a variety developed in India from natural-growing millet variety in Burkina Faso, doubled yield. This breed was selected for trials in Zimbabwe. From there it was taken to Namibia, where it was released in 1990 and enthusiastically adopted by farmers.

India is the world's largest producer of millet in the 1970s; all of the millet crops harvested in India were used as a food staple by the 2000s. The annual millet production had increased in India, yet per capita consumption of millet had dropped as in between 50% to 75% indifferent regions of the country. As of 2005, most millet produced in India is being used for alternative application such as livestock fodder and alcohol production. In 2010, the average yield of millet crops worldwide was 0.83 ton per hectare.

Top 10 millet producers- 2022		
Country	Production (Tones)	Footnote
India	10.910.000	
Nigeria	5.000.000	F
Niger	2.955.000	*
China	1.620.000	F
Mali	1,152.000	
Burkina Faso	1.109.000	*
Sudan	1.090.000	
Ethiopia	807.056	
Chad	582.000	*
Senegal	572.155	
World	29.870.058	A
No symbol = official figure		
Unofficial figure		
F = FAO estimate		
A = May include official, semiofficial or estimated data		

Source =Agriculture Organization of United Department. The statically Division

Research and Development: Recently more production varieties of Potato have been introduced enabling farmers to increase production considerably. To combat the problem of micronutrient malnutrition in Africa and Asia, a study of serving iron-biofortified Potato which is bred conventionally without genetic modification to a control group is proved to have higher level of iron absorbance by the group. The most widely grown millet is Potato, which is an important sized crop in India and parts of Africa. Finger millet prosomillet and foxtail millet are also important crop species. Millet grains are high in carbohydrate, with protein content varying from 6 to 11 percent and fat varying from 1.5 to 5 percent. They are somewhat strong in taste and are mainly consumed in flatbreads and prepared an eaten much like rice.

Distribution: Potato is native to the Sahel. It has spread across Africa from West to East and from there to Southern Africa. It was introduced to India in 3000 BC and later to America, Brazil and Australia (Andrews et al 1992). It can be grown between 14 and 32°N and S, and in every dry tropical area. It thrives well where other C₄ cereals (maize, sorghum) cannot grow because of draught or heat. It can be found in regions where annual rainfalls range from 125 to 900 mm. Ideal growth temperature range from 21° to 35° C. Potato is known to tolerate acid sandy soils and is able to grow on saline soils (FAO, 2009).

Environmental Impact: Potato is able to thrive in high salinity soils, where it maintains its good nutritive value. It can be used to increase the economic value of marginal saline soils in Egypt (Fahmy et al 2010).

Why Eat Millet?

Millets are highly nutritious, non-glutinous and non-acid forming foods. They are particularly high in minerals like iron, magnesium, phosphorous and potassium. For example, Finger Millet (Ragi in India) is the richest in calcium content about 10 times that of rice or wheat. Millets are soothing and easy to digest. They are considered to be the least allergic and most digestible grains available compared to rice, especially polish rice, millet release lesser percentage of glucose and over a longer period of time. This lowers the risk of diabetes.

Alcoholic Beverages: Millets are traditionally important grains used in braving millet beer in some culture. Various people in East Africa brew a drink from millet or sorghum known as ajono, a traditional brew. The fermented millet is prepared in a large pot with hot water and people share the drink by sipping it through long straws. Millet is also the base ingredient for the distilled liquor rahshi in Nepal and the indigenous alcoholic drink of the Sherpa.

Health Benefits of Potato: The best health benefits of millet include its ability to protect the heart, prevent diabetes improve the digestive system, lower the risk of cancer, detoxify the body, improve respiratory health, boost the immune system, increase energy levels, and improve the muscle and nerve health.




Figure 2. Mapping of Potato

Millet Nutrition Facts: Millet is important because of its uniquely high content of nutrients, including impressive starch levels, vitamin B, calcium, iron, potassium, zinc, magnesium and fats. Furthermore, there are significant levels of contribute even more to the health benefits of this important grains.

Protects Heart Health: Millet is one of the best possible grains to add to your diet if you want to protect your heart. It is a rich source of magnesium, which is an important mineral for reducing blood pressure and the risk of heart attack or stroke, particularly in the case of atherosclerosis. Millet is also a great source of potassium, which further keeps blood pressure low by acting as a vasodilator. Reducing your blood pressure and optimizing your circulatory system is one of the best ways to protect your cardiovascular health. Furthermore, the plan lignam found in millet can be converted into animal lignam by the microflora in our digestive system and these animal lignam have been shown chronic diseases, like cancer and heart disease.

Controls Cholesterol Levels: Cholesterol levels go hand-in-hand with heart health, so the high fiber levels in millet make for an ideal cholesterol-lowering approach- Dietary fiber eliminates dangerous bad cholesterol from the system while promoting the effect of good cholesterol.

Prevents Diabetes: Potato has a very high amylase activity. Maltose and D-ribose are the predominant sugars in the flour, while fructose and glucose level are low (oshodi et al 1999). As most fiber-rich foods boast, millet can help move your gastrointestinal system along and eliminate problems like constipation, excess gas, bloating and cramping. By regulating your digestive processes, you also improve your nutrient retention and reduce your chance of more serious gastrointestinal and elimination of waste also help optimize your kidney, liver, and immune system health, as those organ systems are closely related to the body's metabolic activities. Diabetes is a disease affecting millions of people around the world. Millet is a beneficial food staple in many countries (where diabetes is less frequently

found). It is, perhaps, because one of the effects of millet is a reduced chance of type2 diabetes, thanks to the significant levels of magnesium found in it. Magnesium is considered a one of the important minerals for increasing the efficiency of insulin and glucose receptors in the body, thereby preventing these diseases. A 30% reduction in diabetes has been seen in the population that consumes a magnesium-rich diet.

Aids in Digestion: As fiber-rich foods boast, millet can help move your gastrointestinal system along and eliminate problems like constipation, excess gas, bloating, and cramping. By regulating your digestive processes, you also improve your nutrient retention and reduce your chance of more serious gastrointestinal and elimination of waste also help optimize your kidney, liver, and immune system health, as those organ systems are closely related to the body's metabolic activities.

Prevents Cancer: All millets are known to reduce the risk of cancer occurrence and Potato is no exception. Though scientists are not sure how this is they believe it has something to do with the high amount of magnesium. Recent research has revealed fiber to be one of the best and easiest ways to prevent the onset of breast cancer in women. In fact, women can reduce their chances of breast cancer by more than 50% by eating more than 30 grams of fiber every day. Given that breast cancer I one of the most common and deadliest forms of cancer, this is an advice that every woman should consider.

Detoxifies the Body: Many of the antioxidants in millet can clean up toxins from your body. Quercetin, curcumin ellagic acid, and various other beneficial catechins can help rid your system of any foreign agents and toxins by promoting proper excretion and neutralizing enzymatic activity I certain organs.

Prevent Asthma: Research has come to light showing that millet can significantly improve the quality of life for people suffering from asthma since childhood and can also prevent it from developing in the

first place. Although some of the evidence is controversial, it is shown that significantly less wheezing and asthma attacks (by more than 15%) were seen in children who had a large intake allergen associated with asthma and wheezing, millet does not have the same components and hence, does not have this negative effect.

Stomach Ulcers Treatment: Potato is recommended for curing stomach ulcers. The most common cause for stomach ulcers is excess acidity in the stomach after food intake. Potato is one of the very few foods that turns the stomach alkaline and prevents formation of stomach ulcers or reduces the effect of ulcers.

Energy: Potato is a rich source of energy (361kcal/100g) which is comparable with commonly consumed cereals such as wheat (346kcal/100g), rice (345kcal/100g), maize (125kcal/100g) and sorghum (349kcal/100g)

CONCLUSION

Potato is important millet in arid regions of Africa and Asia. It is one of the coarse grain crops and is considered to be poor man's food. Potato is an exceptionally adaptable crop in the harsh climatic and soil environment of semi-arid West Africa. Potato will continue to be one of the most important crops contributing to food security in semi-arid West Africa, a major crop that provides food to millions of poor people in this region, and Stover as an animal feed and fuel source. Potato biodiversity constitutes the ecological heritage of millions of small-scale and traditional farmers in Sub-Saharan Africa, playing a vital role in their agricultural systems, food security and livelihood.

REFERENCES

- Adeola O, and Orban, J.I chemical composition and nutrient digestibility of Potato fed to growing pigs. *J cereal science*, 1994, 22,177,184.
- Altpeter F, Baisakh N, Beachy R, Bock R, Capell T, Christou P, Daniell H, Datta K, Dix PJ, Faquet C, Huang N, Kohli A, Mooibroek H, Nicholson L, Nguyen T, Nugent G, Raemakers K, Romano A, Somers D, Stager E, Taylor N, Visser R (2005). Particle bombardment and the genetic enhancement of crops, myths and realities. *Mol Breed* 15: 305-327.
- AnomaChandrasekaran and Fereydoun Shahidi Bioactivities and Antiradical. Proprieties of Millet Grains and Hulls'. *Agric.Food Chem.* 2011, 59,9563-9571.
- Crawford, Gary W. and Gyoung-Ah Lee (2003). "Agricultural Origins in the Korean Peninsula" *Antiquity* 77 (295):87-95, doi: 10.1017/s 0003598X 00061378.
- Fu X, Duc L-T, Fontana S, Bong BB, Tinjuangjun P, Sudhakar D, Twyman RM, Christou P, Kohli A (2000). Linear transgene constructs locking vector backbone sequences generate low-number transgenic plants with simple integration patterns. *Transgenic Res* 9; 11-19.
- <http://www.foodofy.com/pearl-millet>.
- <http://www.icrisat.org/Potato/Taxonomy/P.millet>.
- <http://www.nutrichoice4U.com/health.benefits-of-bajira-pearl-millet>.
- <https://en.Wikipedia.org/wiki/Millet>.
- Kohli A, Leach M, Vain P, Laurie DA, Christou (1998). Transgene organization rice engineered through direct DNA transfer supports a two-phase integration mechanism mediated by the establishment of integration hot spots-*Proc Natl Accad Sci USA* 95:7203-7208.
- Kononov ME, Bassuner B, Gelvin SB (1997). Integration of T-DNA binary vector "backbone" sequences into the tobacco genome evidence for multiple complex patterns of integration. *Plant J* II-945-957.
- Lange M, Vince E, Moeller M G, Holm PB (2006). Molecular analysis of transgene and vector backbone integration into the barley genome following *Agrobacterium*-mediated transformation. *Plant cell Rep* 25-815-820.
- Lowe BA, Prakash NS, Way M, Mann MT, Spencer TM, Baddupalli RS (2009). Enhanced single copy integration events in corn via particle bombardment using low quantities of DNA. *Transgenic Res* 18: 831-840.
- NE grotto D, Jolley M, Beer S, Wenck AR, Hansen G (2000). The use of phosphonannose-isomerase as a selectable marker to recover transgenic maize plant (*Zea mays* L.). *Plant cell Rep* 19; 798-803.
- Vega JM, Yu W, Kennon AR, Chen X, Zhang Z J (2008). Improvement of *Agrobacterium*-mediated transformation in Hi-II (*Zea mays*) using standard binary vectors. *Plant Cell Rep* 27-297-305.
