



Full Length Research Article

IMPACT OF BODY MASS INDEX ON FASTING BLOOD GLUCOSE IN POSTMENOPAUSAL WOMEN

¹Meenakshi Sharma, ²Shaugfta Aara, ³Anjali Nadir Bhat, ⁴Sushil Sharma and ^{5,*}Shahid Hussain

^{1,2,3}Department of Physiology, Government Medical College, Jammu, Jammu and Kashmir, India

⁴Department of Internal Medicine, Government Medical College, Jammu, Jammu and Kashmir, India

⁵Department of Community Medicine, Government Medical College, Jammu, Jammu and Kashmir, India

ARTICLE INFO

Article History:

Received 11th November, 2016
Received in revised form
14th December, 2016
Accepted 07th January, 2017
Published online 28th February, 2017

Key Words:

Body Mass Index,
Postmenopausal,
Fasting blood glucose

ABSTRACT

Objectives: To assess the relationship of body mass index with fasting blood glucose in postmenopausal women.

Methods: The study was conducted on 100 postmenopausal women in the age group of 45-55 years. Subjects were divided into 3 subgroups normal, overweight and obese on the basis of BMI. Body weight and height was taken and BMI was calculated using Quetlet index. Blood samples were taken in the early morning after a minimum 8 hours of fasting and were investigated for fasting blood glucose. Data obtained was compiled and statistical analysis was done. All parameters were reported as mean and standard deviation. The statistical differences in mean values were tested using Levene' test for equality of variances followed by t-test for equality of means. ANOVA was used to do comparison with in the subgroups. Pearson correlation was used to calculate the correlation between various variables. A p-value <0.05 was considered statistically significant and p-value >0.05 was considered non-significant.

Conclusion: A highly significant correlation was seen between fasting blood glucose and BMI in normal, overweight as well as obese postmenopausal women.

Copyright©2017, Meenakshi Sharma et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Correlation of BMI with blood glucose in women is an important aspect for viewing the impact of changing hormonal levels on the advancing age of women, as menopause sets in. Menopause, a normal biological event marked for most women by the end of menstrual period, signifies the depletion of functional ovarian follicles, that are responsible for estradiol production (Johnson, 1998). Menopause is the permanent cessation of menstruation resulting from the loss of ovarian follicular activity. It is heralded by the menopausal transition, a period when the endocrine, biological and clinical features of approaching menopause begin. A common initial marker is the onset of menstrual irregularity. The biology underlying the transition to menopause includes central neuro-endocrine changes as well as changes within the ovary, the most striking of which is a profound decline in follicle no.'s (Burger *et al.*, 2002) Menopause is associated with energy expenditure during rest and physical activity, an accelerated loss of fat-free mass and alteration of adipose tissue metabolism and fat oxidation. This deregulation of energy metabolism could induce an increase in total adiposity and redistribution of fat in

the abdominal region in postmenopausal women (Paszkowski and Klodnicka, 2007; Funahashi *et al.*, 2004). As BMI and central obesity pose a risk for diabetes mellitus, assessment of correlation of BMI with blood glucose and postmenopausal women is essential as obesity is the most common disorder associated with women in their menopausal stage and occurs in approximately 65% of women (Koerner *et al.*, 2005). Obesity is associated with risk of hypertension, atherosclerosis, dyslipidemia, cancer, osteoarthritis and respiratory problems (Fulam *et al.*, 2006; Ferrara *et al.*, 2002; Kimura *et al.*, 2006). So, the present study is planned to assess the effects of increase in bodyweight on fasting blood glucose in women who have attained menopause. It can help the women of advancing age to pay heed to the modifiable risk factors so that overall morbidity due to diabetes can be decreased in these women.

MATERIALS AND METHODS

Selection procedure of the subjects

Subjects were selected from different outpatient departments of Government Medical College, Jammu. Subjects included healthy postmenopausal women in the age group of 45 to 55 years.

*Corresponding author: Shahid Hussain

Department of Community Medicine, Government Medical College, Jammu, Jammu and Kashmir, India.

Table 1. Mean BMI in subgroups of postmenopausal women

BMI (Kg/ m ²)	Classification	Postmenopausal (no.) Group II	Mean BMI ± SD (Range) (in Kg/ m ²)
18.50 – 24.99	Normal weight	37	21.97± 1.12 (18.9 – 24.03)
25.00 – 29.99	Overweight	34	27.61 ± 1.42 (25.14 – 29.74)
≥ 30.00	Obese	29	32.55 ± 2.41 (30.04 – 37.78)

Table 2. Correlation of BMI with fasting blood glucose (FBG) in postmenopausal women

BMI (Kg/ m ²)	Classification	Postmenopausal Women (no.) Group II	Mean fasting Blood glucose ± SD(Range)	Pearson correlation with p value
18.50 24.99	Normal weight	37	99.64+13.7 (70-130)	0.432 P=0.008
25.00 29.99	Overweight	34	97.38+9.98 (86-121)	0.818 P=0.000
> 30.00	Obese	29	108.75+18.74 (72—150)	0.866 p =0.00

Table 3. Comparison of fasting blood glucose (FBG) and BMI in postmenopausal group

Parameter	BMI			F Value	p Value
	Normal weight Mean ± SD (n=37)	Overweight Mean ± SD (n=34)	Obese Mean ± SD (n=29)		
FBG	94.64±13.70	97.38±9.98	108.75±18.74	8.532	0.000

All those women with H/O diabetes mellitus, hypertension, neurological disorders, any other illness known to affect fasting blood glucose, were excluded from the study, as were smokers and alcoholics. For body mass index, body weight was measured in kilograms, on a weighing scale (Stranges *et al.*, 2004). Height in meter was also measured. Body mass index was calculated using Quetelet's Index (Pranita *et al.*, 2012).

Specimen collection and handling for estimation of blood glucose

A fasting blood sample of 2 ml was drawn with appropriate aseptic precautions in the early morning after a minimum of 8 hours fasting. The glucose method is an adaption of the hexokinase-glucose-6-phosphate dehydrogenase method. Subjects were further divided into 3 subgroups namely normal, overweight and obese.

Statistical Analysis

The statistical differences in mean values were tested using Levene's test for equality of variances followed by t-test for equality of means. ANOVA was used to do comparison with in the subgroups. Pearson correlation was used to calculate correlation between various variables.

RESULTS

Postmenopausal women

The BMI range in normal weight females is 18.9 to 24.03 Kg/m², with a mean of 21.97 (SD ± 1.12) Kg/m². Overweight females have BMI range of 25.14 to 29.74 Kg/m², with a mean of 27.61 (SD ± 1.42) Kg/m². Obese females have BMI range of 30.04 to 37.78 Kg/m², with a mean of 32.55 (SD ± 2.41) Kg/m². Fasting blood glucose range in normal weight females is 70-130mg/dl, with a mean of 94.64 (SD+ 13.7)mg/dl.

Overweight females have fasting blood glucose range of 86 to 121 mg/dl, with a mean of 97.38 (SD +9.98) mg/dl. Obese females have fasting blood glucose range of 72 to 150 mg/dl, with a mean of 108.75 (SD+18.74) mg/dl. There is a positive correlation in all 3 subgroups and p- value is highly significant in all the 3 subgroups. Post menopausal women show significant difference of fasting blood glucose (FBG) in all three subgroups and p-value is highly significant. There is increase in blood glucose in correspondence with increase in BMI in postmenopausal women.

DISCUSSION

The abrupt endocrine changes during menopausal transition have important impacts on the physiology of female body which exacerbates risks for many diseases and disabilities during postmenopausal life (Mishra, 2011), mainly estrogen deficiency. Postmenopausal status is associated with a 60% increased risk for metabolic syndrome. It is thought to be associated with decreased estrogens and increased abdominal obesity in postmenopausal women with metabolic syndrome (Koh *et al.*, 2008). Increase in BMI lead to increased levels of blood glucose, hypertension, dyslipidemia, metabolic syndrome. Our study showed that BMI has high impact on the fasting blood glucose in postmenopausal women. Need *et al.* 2005 also observed in their study that fasting serum glucose was positively and significantly correlated with BMI. The findings support our observations that BMI as well as central obesity pose a risk factor for diabetes. BMI was greater for women who had just undergone menopause (Ueda *et al.*, 1995). Mean blood glucose level in the menopausal women increased slightly than that of the non-menopausal women (Duru *et al.*, 2014). Koskova *et al.* (2009) observed in their study that fasting serum glucose increased with age, increasing slightly up to menopause and the postmenopausal group showed a significant increase with frequent pathological values. C- peptide increased slightly with age. Increase in insulin was evident up to menopause. Increase in fasting glucose level is greater in postmenopausal women (Matthews

et al., 2001). Many postmenopausal women gain weight, which leads to greater incidence of type 2 diabetes (Reckelhoff and Fortepiani, 2004). Obesity is related to ectopic accumulation of lipids in the muscles, liver and beta-pancreatic cells, leading to insulin resistance in the muscles and liver and impairment of beta- cells function. Free fatty acids are in competition with glucose for oxidation, suggesting that increased lipid oxidation worsens insulin resistance in obesity (Savage et al., 2005). This study was also in agreement with the fact that menopause results in a decrease in both pancreatic insulin secretion and insulin elimination. A larger percentage of type 2 diabetics have central obesity as compared to general obesity and both types of obesity was greater in females as was observed by Kamath et al.(2011). Assessment of body weight composition in postmenopausal women and determination of correlations with metabolic and hormonal parameters was done. The authors found that body mass index values positively correlated with age, time since menopause, parity and glucose, thereby displaying significant correlations with hormonal and metabolic parameters (Cuadros et al., 2011).

Conclusion

We conclude that highly significant correlation was seen between fasting blood glucose and BMI in normal, overweight as well as obese postmenopausal women.

REFERENCES

- Johnson, S.R. 1998. Women's health issues. *Med Clin North Am.*, 82:
- Burger, H.G., Dudley, E.C., Robertson, D.M. and Dennerstein, L. 2002. Hormonal changes in the menopause transition. *Recent Prog Horm Res.*, 57: 257-275.
- Paszkowski, T. and Klodnicka, M. 2007. Hormonal therapy of menopause. *Menopause*, 2: 106-109.
- Funahashi, T., Matsuzawa, Y. and Kihara, S. 2004. Adiponectin as a potential key player in metabolic syndrome. Insights into atherosclerosis, diabetes and cancer. *Int Congress Sci.*, 1262: 368-371.
- Koerner, A., Kratzch, J. and Kiess, W. 2005. Adipocytokines: leptin – the classical resistin – the controversial adiponectin. The promising and more to come. *Best Pract Res Clin Endocrinol Metab.*, 19: 525-546.
- Fulam, T., Tessier, D. and Carpentier, A. 2006. The metabolic syndrome. *Pathol Biol.*, 54: 375-386.
- Ferrara, C.M., Lynch, N.A., Nicklas, B.J., Ryan, A.S. and Berman, D.M. 2002. Difference in adipose tissue metabolism between postmenopausal and perimenopausal women. *J Clin Endocrinol Metab.*, 87: 4166-4170.
- Kimura, T., Matsumoto, T., Akiyoshi, M., Owa, Y., Miyaska, N. and Aso, T. 2006. Body fat and plasma lipids in postmenopausal women are related to resting autonomic nervous system activity. *Eur J Appl Physiol.*, 97: 542-547.
- Stranges, S., Dorn, J.M., Muti, P., Freudenheim, J.L., Farinaro, E. and Russell, M. 2004. Body fat distribution, relative weight, and liver enzyme levels: a population-based study. *Hepatology*, 39(3): 754-763.
- Pranita, A., Phadke, A.V., Singh, R., Melinkeri, R.R. and Joshi, A.R. 2012. Prediabetes status in obese and preobese women in the age group 45-49 years. *J Assoc Phys India*, 60: 121-127.
- Mishra, S.K. 2011. Menopausal transition and postmenopausal health problems: a review on its bio-cultural perspectives. *Health*, 3(4): 233-237.
- Koh, J.H., Lee, M.Y., Nam, S.M., Sung, J.K., Jung, P.M., Noh, J.K., Shin, J.Y., Shin, Y.G. and Chung, C.H. 2008. Relationship between menopausal status and metabolic syndrome components in Korean women. *Korean Diabetes J.*, 32: 243-251.
- Need, A.G., O'Loughlin, P.D., Horowitz, M. and Nordin, CCBE. 2005. Relationship between fasting serum glucose, age, BMI and serum 25-hydroxy vitamin D in postmenopausal women. *Clinical Endocrinology*, 62(6): 738-741.
- Ueda, M., Mizushima, S., Biosca, M. and Sanchez, R.G. 1995. Impact of menopause status on blood pressure: The WHO Cardiac study. *Cardiovasc Risk Factors*, 5: 290-298.
- Duru, B.N., Omonayin, O.J., Gadzama, H.M., Olawuyi, N.O., Diddam, M.A., Nambol, D.L., Ujah, A.E., Ashi, R.R., Okoye, N.O., Tyem, D.D., Udom, I.E., Salako, C.Y., Adeyanju, O.N., Gushe, D.D., Abilu, C.J. and Danjuma, L.A. 2014. Assessment of fasting blood plasma glucose level in menopausal women in rural area of Vom and Environs. *IOSR-JNHS*, 3(3): 57-60.
- Koskova, I., Petrsek, R., Vondra, K., Duskova, M. and Starka, L. 2009. Metabolic profile and sex hormone binding globulin in different reproductive phases of Czech women and their relation to weight, body composition and fat distribution. *J Physiol.*, 38 (2): 393-402.
- Matthews, K.A., Kuller, L.H., Sutton-Tyrrell, K. and Chang, Y.F. 2001. Changes in cardiovascular risk factors during perimenopause and postmenopause and carotid artery atherosclerosis in healthy women. *Stroke*, 32: 1104-1111.
- Reckelhoff, J. and Fortepiani, A. 2004. Novel mechanisms responsible for postmenopausal hypertension. *Hypertension*, 43: 918-923.
- Savage, D.B., Petersen, K.F. and Shulman, G.I. 2005. Mechanisms of insulin resistance in humans and possible links with inflammation. *Hypertension*, 45: 828-833.
- Kamath, A., Shivaprakash, G. and Adhikari, P. 2011. Body mass index and waist circumference in type 2 diabetes mellitus patients attending a diabetes clinic. *Int J Biol Med Res.*, 2(3): 636-638.
- Cuadros, J.L., Fernandez-Alonso, A.M., Cuadros, A.M., Chedraui, P. and Perez-Lopez, F.R. 2011. Body mass index and its correlation to metabolic and hormonal parameters in postmenopausal Spanish women. *Gynecol Endocrinol.*, 27(9): 678-684.
