

Available online at http://www.journalijdr.com



International Journal of Development Research Vol. 08, Issue, 02, pp.18798-18801, February, 2018

ORIGINAL RESEARCH ARTICLE



OPEN ACCESS

MOTOR FITNESS AMONG INDIAN AND IRANIAN STUDENTS A- CROSS SECTIONAL STUDY

*Majid Hashemi

Assistant Professor in Department of Physical Education, Varamin-Pishva branch, Islamic Azad University, Varamin, Iran

ARTICLE INFO

Article History: Received 20th November, 2017 Received in revised form 13th December, 2017 Accepted 19th January, 2018 Published online 28th February, 2018

Key Words:

Motor fitness, Motor Behavior, Movement, Students' age, leg power, Cross-Sectional Study.

ABSTRACT

This paper aims to compare the motor development of Indian and Iranian boy's student between the age of 13 and 18 years in explosive leg power case. The focus is on the relationship between age and motor behavior which makes the study of motor development unique from other viewpoints. Motor development includes age related changes in both posture and movement, the two basic ingredient of motor behavior. The results portrayed by means of statistical tests and standard method of sampling.

Copyright © 2018, *Majid Hashemi*. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Majid Hashemi. 2018. "Motor fitness among indian and iranian students a- cross sectional study", International Journal of Development Research, 8, (02), 18798-18801.

INTRODUCTION

This study was conducted keeping in view the following objectives. Study the level and pattern of the development of motor abilities of Indian male student of 13-18 years' age. Compare the level and pattern of development of motor abilities of Indian male student with Iranian male student.

Definitions of operational terms motor development

Motor development is the process of change in motor behavior that is related to the age of the individual. The focus on the relationship between age and motor behavior makes the study of motor development unique from other viewpoints motor development include age related changes in both posture and movement, the two basic ingredient of motor behavior. Development processes occur throughout the human life span (Jan Stephen tecklin, 1998). For the purpose of this study the term was understood to mean motor ability through the performance in selected motor fitness component that underlie gross motor skills.

Corresponding author: Majid Hashemi,

Assistant Professor in Department of Physical Education, Varamin-Pishva branch, Islamic Azad University, Varamin, Iran.

Cross-Sectional Study

The cross-sectional study is a method of study that permits the researcher to collect data on different groups of people at varying age levels at the same point in time. The major purpose of the cross-sectional study was to measure of age-related difference in behavior. This method does not permit measurement of age-related change, and has attracted controversy in recent years. Basically the cross-sectional method yields only average difference in groups across real time and not individual change developmental time. The basic assumption behind the cross-sectional study had been that random selection of subjects provides as represented sample of the population for each age group test. (David L & Gallahue, 1995) p10

Delimitations

Motor development is an all-inclusive which is the area of interest for child psychologists, social psychologists and sport psychologists alike. Thus the term motor development is much wider in scope and meaning. For the purpose of this study the term motor development was contained to the concept of motor fitness development as measured through recognized motor fitness component of strength. The study was also delimited to high school student of 13-18 years. The study was further delimited to Indian students in Chandigarh and Iranian student in Tehran. The study was further more delimited to male students.

Limitation

Even though, no motivational techniques were employed, but every effort was made by the researcher to encourage the subjects to do their best. In spite of that researcher could not possibly control the performance differences in effort made by the subject to do their best. Such variation in effort and home environment, daily routine and diet might distort actual scores collected through and ultimately the final analysis. Variations obtained in score due to this factor were duly recognized as the limitation of the study.

Significance of the study

In the past one-decade physical education has found its right place in the school curriculum. To a large extent it has found its academic recognition at par with other subjects. Having found its place in the school curriculum, the teacher of physical education is confronted with numerous problems relating to classroom instruction in physical education. One specific problem that relates to instruction is the extent to which the school student may be provided combine instruction irrespective of their age. Another problem is catering to the individual needs. Even though providing the individualized instruction in physical education may be a far off dream, yet the teacher had to ensure that each group has only acceptable variation in abilities in order to provide effective instruction and avoid damages. The teaching policies signs and teachers, there for, should be well familiar with the development trends and generalized pattern of development at different stages. This may help to adjust to programmers to the needs of the group as a whole. Thus, the present investigation may be great significance in understanding the patterns of development in motor ability. This may help to draw out effective learning environment and to provide scope for individual attention to extent possible. The result of the study may also help to understand the classification criterion in a better way. The result of the study helps to understand the role of diet patterns, topography, genetic factor and the effect of educational system on school going children, in affecting the development pattern of children.

Literature

Haley (1972) conducted a study of motor fitness. The sample included children studying in grades one through six. Thirty boys were randomly selected from each grade. Their ages ranged from five years nine months to 12 years two months. Sharma and others (2017) Studied Anthropometric Basis of Vertical Jump Performance in Young Indian National hockey and cycling players of 11-21 years in variables: Weight, Height, Trochanter Ion-Height, Sum of Skinfold Thickness, Lengths [Acromiale-Stylion and Midstylion-Dactylion], Breadths [Biacromial, Biiliocristal, BiepicondylarHumerus and Biepicondylar Femur], Girths [Relaxed Arm, Mid-Thigh and Calf], Lower Back Strength. Schmidt (1982) after reviewing numerous studies pointed out which by the time an individual researches the age of 18 he experiences large improvement in his motor behavior. HaiphotChanchiclung (1985) conducted an assessment of physical fitness of lower secondary school boys of Thailand. The samples for the study

were 13500 lower secondary school boys selected through randomized clustered sampling. The modified Fleishman physical fitness test battery which consists of item for flexibility, quickness, strength, muscular endurance, matching, balance and cardiovascular endurance was administrated to the Reet Mahindersingh (1986) Pr epared physical subjects. fitness norms for high school boys of panjab state. Data were collected on 5000 subjects selected randomly from various schools in the state. The test battery managed comprised of eight items. I.e. Rachpal Singh Brar (1987). Conducted in effects of short interval and long interval running with two recovery types on aerobic and anaerobic capacities and running performance of high school boys, the subject were 100 untrained students of grades nine and ten in Shivalik public school. DaljitKaur (1989). Conducted a study on the physical fitness of high school girls of the panjab belonging in the age group of 12 to 15, the purpose of the study was to prepare norms for the girls of panjab belonging to this age group. MeeraChauhan (1989). Compared the motor fitness performance of sports and non-sports school girls (1315) years' old living at the high altitude of 2960ms at Shimla and 487ms at Chandigarh, SukhpalKaur (1990). Conducted acrosssectional study of motor abilities of panjab and Chandigarh girls in the age group of7 to 11 years, the investigator studied the developmental changes in motor abilities which take place during the mentioned period.

Amarpreet Singh (1993) conducted a study on the relationship of varying levels of motor fitness to Socio-Economic statues and structural variations among school students in the age group of 14 to 16 years. Shilendra Kumar Sinha (1996) conducted a study of anthropometric and motor quality profiles of 8-14 years' boys of eastern and north east region of India. Kamal Kant Sharma (1997) conducted a study on construction and standardization of motor fitness test battery for elementary school children in Delhi (U.T), the objectives of study were as follow: to find out how motor fitness variables, such as speed, strength, balance, flexibility and endurance, develop among boys and girls in the age group of eight to eleven years. Dinesh kumar (1998) showed on a normative study of fitness status in male students (13-16) years of age belonging to the schools of Himachal Pradesh, followed by development of norms for future uses. JasbireKaur (1999) Conducted as assessment of motor fitness of rural and urban senior secondary school girls of Punjab state. SonamAngchok (1999) conducting a study to establish norms for the high and higher secondary male student of ladakh, among the age group 13 to 17. Sujata Devi (2000) conducted a study to compare the physical fitness and psychological trait of tribal and non-tribal high school students of high altitude areas between the age group of 14 and 17 years. MandeepBrar (2004) conducted a study on motor development of school children of union territory of Chandigarh a cross sectional analysis 12 to 14 vears.

MATERIALS AND METHODS

In this chapter selection of subjects, design of the study, selection of variables, reliability of data, tools used, reliability of instrument, criterion measure, collection of data, administration of selected test items for collection of data, and techniques for data analysis are described.

Selection of subjects

The selection of subjects was completed in two phase 1 - a pilot study had been conducted on 240 student of 13-18 years

Table 1. The analysis of variance of Indian and Iranian male students in Standing Broad Jump

Source of variation	DF	S	S	M	S		F
		Indian	Iranian	Indian	Iranian	Indian	Iranian
Between groups	5	107600.794	142546.574	21520.159	28509.315	46.080*	44.732*
Within groups	1194	557619.305	767153.905	467.018	642.507		
Total	1199	665220.099	909700.479				

*Significant at .05 Level of Confidence.

of age studying in government schools from classes seven to twelve, 120 from Chandigarh (India) and 120 from region nine of Tehran (Iran) 20 students from each age, Abbreviations GSSS, GHSSS, GMSSS and JNVS means: government senior secondary school, government high school senior secondary, government model senior secondary school and Jawahar Navodaya Samiti respectively.

Collection of data

The data for selected variables on the randomly selected subjects was collected over a period of eleven months (12/01/09 to 02/12/09). The subjects were made available by school authorities during the physical education classes and other times when the students were available from their regular academic routine. So the data was collected over different times of the day for different variables.

Statistical technique employed

To establish the reliability of the data person product moment correlation method was used. In order to analyze development patterns in motor fitness, analysis of variance was carried out for motor fitness item to determine significance of variance, if any, from age to age, separately for Indian and Iranian students. Whenever F values were found significant, the posthoc scheffe's test was employed to determine the significance of difference between the paired means. For analyzing difference between Indian and Iranian at each age in motor fitness test item, the t test is applied. The level of significant was set at .05. The table 1 showed a significant F values of 46. 080 for Indian and 44.372 for Iranian student respectively, which indicated that six age groups different significantly in strength. Because the F was found to be significant, to establish which paired age group differed the results of post hoc scheffe's test have been present in tables 2-7.

Table 2. comparison of the paired mean test means for respective categories of Indian and Iranian male students in Standing Broad Jump

GROUPS	MD		P value	
	Indian	Iran	Indian	Iran
13 vs. 14	9.150*	15.220*	.003	.000
13 vs. 15	14.565*	25.155*	.000	.000
13 vs. 16	21.215*	26.555*	.000	.000
13 vs. 17	25.890*	33.720*	.000	.000
13 vs. 18	26.405*	26.345*	.000	.000

*Significant at .05 level of confidence.

It was evident from the table that in Indian male students paired mean differences

Table 3. comparison of the paired mean test means for respective categories of Indian and Iranian male students in Standing Broad Jump

		P		
GROUPS	MD		P value	
	Indian	Iran	Indian	Iran
14 vs. 13	9.150*	15.220*	.003	.000
14 vs. 15	5.415*	9.935*	.281	.009
14 vs. 16	12.065*	11.335*	.000	.001
14 vs. 17	16.740*	18.500*	.000	.000
14vs. 18	17.255*	11.125*	.000	.002

*Significant at .05 level of confidence.

It was evident from the table that in Indian male students paired mean differences

It is evident from the table 1, in Indian male students paired mean difference of 13 and 14years, 13 and 15years, 13 and 16years, 13 and 17 years, and 13 and 18 years are significant respectively.

Table 4. comparison of the paired mean test means for respective categories of Indian and Iranian male students in Standing Broad Jump

GROUPS	MD		P value	
	Indian	Iran	Indian	Iran
15 vs. 13	14.565*	25.155*	.000	.000
15 vs. 14	5.415	9.935*	.281	.009
15 vs. 16	6.650	1.400	.093	.998
15 vs. 17	11.325*	8.565*	.000	.044
15 vs. 18	11.840*	1.190	.000	.000

*Significant at .05 level of confidence.

Table 5. comparison of the paired mean test means for respective categories of Indian and Iranian male students in Standing Broad Jump

GROUPS	MD		P value	
	Indian	Iran	Indian	Iran
16 vs. 13	21.215*	25.555*	.000	.000
16 vs. 14	12.065*	11.335*	.000	.001
16 vs. 15	6.650	1.400	.093	.998
16 vs. 17	4.675	7.165	.457	.158
16 vs. 18	5.190	.210	.330	1.000

*Significant at .05 level of confidence.

This indicated that in broad jump 14, 15,16, 17 and 18-year male student were belter than 13 year male students. In Iranian male students the mean differences of 13 and 14 years, 13 and 15 years, 13 and 16 years, 13 and 17 years, and 13 and 18 years are significant respectively. This indicated that in broad jump 14, 15, 16, 17 and 18-year male student were belter than 13 year male students. From the table 2, it is evident that in Indian male student paired mean difference of 14 and 13 years, 14 and 16 years, 14 and 17 years, and 14 and 18 years were found to be significant respectively. This indicated that in strength 13, 16, 17 and 18-year male student were better than 14 year male students, also indicated that 13 year was not better than 14 years and 14 years' male students was not better than 16, 17 and 18 years' male students.

Table 6. comparison of the paired mean test means for respective categories of Indian and Iranian male students in Standing Broad Jump

GROUPS	MD		P value	
	Indian	Iran	Indian	Iran
17 vs. 13	25.890*	33.720*	.000	.012
17 vs. 14	16.740*	18.500*	.000	.000
17 vs. 15	11.325*	8.565*	.000	.044
17 vs. 16	4.675	7.165	.457	.158
17 vs. 18	.515	7.375	1.000	.133

*Significant at .05 level of confidence.

Table 7. comparison of the paired mean test means for respective categories of Indian and Iranian male students in Standing Broad Jump

GROUPS	MD		P value	
	Indian	Iran	Indian	Iran
18 vs. 13	26.405*	26.345*	.000	.000
18 vs. 14	17.255*	11.125*	.000	.002
18 vs. 15	11.840*	1.190	.000	.999
18 vs. 16	5.190	.210	.330	1.000
18 vs. 17	.515	7.375	1.000	.133

*Significant at .05 level of confidence.

In Iranian male students the paired mean difference of 14 and 15, 16, 17 and 18 years was significant this indicated that 14 year was better than 13 years of age. From the table 3 It was evident that in Indian male students paired mean differences of 15 and 13 years, 15 and 17 years and 15 and 18 years were found significant. This indicated that in strength performance 15 year was better than 13 years and 17 and 18 years was better than 15 years of age. But in cases of 15 and 14 years, 15 and 14 and 15 and 16years were not found significant differences. This indicated that in strength performance 14 year was not better than 15 years and 15 years was not better than 16 years of age. In Iranian male students paired mean differences of 15 and 13 years, 15 and 14 years, 15 and 17 years was found significant. This revealed that in strength performance 15year male students were better than 13 and 14 years. But in cases of 15 and 16 and 18 years of age were not found significant differences. This indicated that in strength performance 15 year was not better than 16, and 18 year male students. From the table6 it was evident that in Indian students between age of 16 and 13, 16 and 14 was found significant, it indicated that 16 years of age was significantly better than 13 and 14 years but in other age group namely 15,17 and 18 was not found significant. In Iranian students the ages of 13 and 14 years was found to be significant, it reveals that 16 years was better than 13 and 14 years of age, but in ages of 15, 17 and 18 years of age was not found to be significant. In table 6 it is evidence that in Indian students between age groups of 17 and 13, 14 and 15 years of age was significant, it reveals that 17 years' students were better than 13, 14 and 15 years of age, but in other age groups namely 16 and 18 years of age was not found significant. In Iranian male students the age groups of 17 and 13, 14, 15 years of age was significant it reveals that 17 years of age was better than 13, 14 1nd 16 years of age, the table 7 reveals that in Indian male students between 18 years' age group and 13, 14 and 15 years of age was significant differences, it indicated that 18 years of age was better than 13, 14 and 15 years of age in strength, also in Iranian male students 18 years of age significantly was better than 13 and 14 years of age.

Conclusion

This paper aims to compare the motor development of Indian and Iranian boy's student between the age of 13 and 18 years in strength performance. The focus on the relationship between age and motor behavior makes the study of motor development unique from other viewpoints. The results and tests showed that six age groups are different significantly in strength. Also the tests indicated that in strength 14, 15, 16, 17 and 18-year male student were better than 13 year male students. In Iranian male students the mean differences of 13 and group 14 and 13, pair 15and 13, pair 16 and 13 and 17 and 13 and 18-year male student were found to be significant. This indicated that in strength 14,15,16,17 and 18 year male students were better than 13 year male students. In final results it derives that Indian male student had significant difference of strength in case of 13 and 15 years, 14 and 16 years, 15 and 17 years, but no significant difference was found in case of 16 and 18 years, 17 years of age was better than other age groups, for the future study the other components like flexibility and etc. proposed to study by the authors.

REFERENCES

- Amarpreet Singh, 1993. "Relationship of varying levels of motor fitness to socio-economic statues and structural variation among school students in the age group of 14-16 years.
- DaljitKaur, 1989. "assessment of physical fitness of high school girls of Panjab".
- David L. Gallahue, John C. Ozmun, 1995. "understanding moor development" (USA WCB Brown & Bench Marck): P10
- Dinesh kumar, 1998. "a normative study of physical fitness of lower secondary school boys (12-14 years of age) of Thailand"
- HaiphotChanchiclung, 1985. "Assessment of motor physical fitness of lower secondary school boys (12-14 years of age) of Thailand"
- JasbireKaur 1999. "Assessment of motor fitness of rural and urban senior secondary school girls of Panjab state".
- Mandeep Brar, 2004. "Motor development of school children of union territory of Chandigarh-A cross sectional analysis 11-14 years". Panjab University Theses.
- Meera Chauhan, 1989. "construction and standardization of motor fitness test battery for elementary school children in Delhi (u.t)
- Philips Roy Haley, "A Comparative Analysis of Selected Motor Fitness Test Performance of Elementary School Boys," Dissertation abstracts International (March 1972): 5018-A
- Rachhpal Singh Brar, 1987. "Effect of short invertal and long interval running with two recovery types on aerobic and anaerobic capacities and running performances of high school boys".
- Reet Mahindersingh, 1986. "Physical fitness norms of Panjab state high school boys".
- Schmidt, R. A. 1982c. The schema concept. In J. A. S. Kelso (Ed.), Human motor behavior: An introduction (pp. 219-235). Hillsdale, NJ: Erlbaum
- Sharma HB, Gandhi S, Meitei KK, Dvivedi J, Dvivedi S, J ClinDiagn Res. 2017 Feb, "Anthropometric Basis of Vertical Jump Performance: A Study in Young Indian National Players". 11(2):CC01-CC05. doi: 10.7860/ JCDR/201 7/23497.9290. Epub 2017 Feb 1.PMID: 28384855
- Shilendra Kumar Sinha 1996. "A study of anthropometric and motor quality profiles of 8-14 years' boys of eastern and north east region of India" kalyani university –SAI, Calcutta.
- Sonam Angchok 1999. "physical fitness norms for high and higher secondary student of Ladakh"
- Sujata Devi 2000. A comparative study of physical fitness and psychological traits of Tribal and non-Tribal high school students of high altitude area".
- Sukhpal Kaur 1990." A cross-sectional study of motor abilities of panjab and Chandigarh girls in the age group of 7 to 11 years". Panjab University Theses.
- Sukhpal Kaur 1990. "A cross-sectional study of motor abilities of girls in the age group of 7 to 11 years".